

Rhodora

JOURNAL OF THE
NEW ENGLAND BOTANICAL CLUB

Conducted and published for the Club, by

BENJAMIN LINCOLN ROBINSON, Editor-in-Chief.

FRANK SHIPLEY COLLINS

MERRITT LYNDON FERNALD

HOLLIS WEBSTER

} Associate Editors.

WILLIAM PENN RICH

EDWARD LOTHROP RAND

} Publication Committee.

Vol. 22.

April, 1920

No. 256.

CONTENTS:

Eupatorium purpureum and its Allies. *K. M. Wiegand* . . . 57

The American *Ammophila*. *M. L. Fernald* 70

Reports on the Flora of the Boston District,—XXXII.

C. H. Knowlton and Walter Deane 72

Juncus Gerardi, var. *pedicellatus*, n. var. *M. L. Fernald* . . . 76

Boston, Mass.

1052 Exchange Building



Providence, R. I.

Preston and Rounds Co.

RHODORA.—A monthly journal of botany, devoted primarily to the flora of New England. Price, \$2.00 per year, postpaid (domestic and foreign); single copies (if available) 20 cents. Volumes 1-8 or single numbers from them can be supplied at somewhat advanced prices which will be furnished on application. Notes and short scientific papers, relating directly or indirectly to the plants of the northeastern states, will be gladly received and published to the extent that the limited space of the journal permits. Forms will be closed five weeks in advance of publication. Authors (of more than one page of print) will receive 25 copies of the issue in which their contributions appear. Extracted reprints, if ordered in advance, will be furnished at cost.

Address manuscripts and proofs to

B. L. ROBINSON, 3 Clement Circle, Cambridge, Mass.

Subscriptions, advertisements, and business communications to

W. P. RICH, 300 Massachusetts Avenue, Boston, Mass.

Single copies may be had from

E. L. RAND, Corresponding Sec'y N. E. Botanical Club,

1052 Exchange Building, Boston, Mass.

Entered at Boston, Mass., Post Office as Second Class Mail Matter.

KEY TO NEW ENGLAND TREES, Wild and Commonly Cultivated, based primarily upon leaf characters, by J. FRANKLIN COLLINS and HOWARD W. PRESTON. Price 40c. net. PRESTON & ROUNDS Co., Providence, R. I.

CARD-INDEX OF NEW GENERA, SPECIES AND VARIETIES OF AMERICAN PLANTS, 1885 TO DATE.

For American taxonomists and all students of American plants the most important supplement to the Index Kewensis, this catalogue in several ways exceeds the latter work in detail, since it lists not only the flowering plants, but pteridophytes and cellular cryptogams, and includes not merely genera and species, but likewise subspecies, varieties and forms. A work of reference invaluable for larger herbaria, leading libraries, academies of sciences, and other centers of botanical activity. Issued quarterly, at \$15.00 per 1000 cards.

GRAY HERBARIUM of Harvard University,

Cambridge, Mass., U. S. A.

CHECK LIST OF GRAY'S MANUAL, 7th EDITION, compiled by M. A. DAY. Leatherette. Pocket size. Invaluable for collector's memoranda and herbarium records. Published and sold by the GRAY HERBARIUM, Cambridge, Mass. Price postpaid 20 cts. each. Ten copies \$1.50.

MEMOIRS OF THE GRAY HERBARIUM. A series of illustrated quarto papers issued at irregular intervals, sold separately.

Vol. I. Monograph of the genus *Brickellia*, by B. L. Robinson (with 96 text figures effectively illustrating all known species of the genus). Feb. 1917. \$3.00.

Gray Herbarium of Harvard University, Cambridge, Mass.

Advertisements of Nurserymen and Dealers in Botanical and other Scientific Publications are inserted in these pages at the following rates per space of 4 in. by 3-4 in. 1 year \$4.00, 6 months \$2.50.

Rhodora

JOURNAL OF

THE NEW ENGLAND BOTANICAL CLUB

Vol. 22.

April, 1920.

No. 256.

EUPATORIUM PURPUREUM AND ITS ALLIES.

K. M. WIEGAND.

THE verticillate-leaved Eupatoriums have long given trouble to American botanists. In connection with work on the flora of Central New York, the writer has found it necessary to make some disposition of the material ordinarily recognized as *Eupatorium purpureum*. That more than one form is to be found under this name in New York State has been apparent for some time, but the limits and characters of these forms did not seem to agree with those recognized by various authors as delimiting the numerous subdivisions of this species. The problem was taken, with others, to the Gray Herbarium, where Dr. Robinson kindly placed at the writer's disposal the very complete collection of *E. purpureum* in that institution, and also photographs made by him of Linnean and Willdenovian specimens in European herbaria. This material was supplemented by the rich collection of New England Eupatoriums in the herbarium of the New England Botanical Club, and of New York State plants of this group in the herbarium of the New York State College of Agriculture. As a result of this study, a treatment has been worked out which it is hoped will aid in clearing up the difficulties that have so long surrounded this section of Eupatorium.

Briefly stated, four species have been confused under the general name *E. purpureum* L. For clearness, these may be numbered, and their most striking characters listed, as follows:

1. Florets 6-9, rarely 5-12: stem speckled: leaves usually in 3's or 4's, ovate, abruptly contracted at the base, more or less plainly

- 3-nerved. Along the Coastal Plain from eastern Massachusetts and southern New Hampshire to South Carolina. A plant of wet soil.
2. Florets 9-15, rarely 8-20: stem speckled: leaves in 4's or 5's, rarely in 3's or 6's, gradually tapering at the base, and pinnately-veined. Newfoundland through northern New England to western Connecticut and central Pennsylvania, westward to Illinois, and Colorado, New Mexico and British Columbia. A plant of wet soil.
 3. Florets 6-7, rarely 5 or 8: stem rarely speckled, evenly purple, glaucous, hollow: leaves in 4's-6's, rather bluntly and finely toothed: corolla less than 5 mm. long. Southern Maine and Rhode Island to Florida, Texas, and Oklahoma; also in western Pennsylvania and Ohio. A plant of damp woods and pastures, on the Atlantic Coast and Uplands.
 4. Florets 5-6, rarely 3-7: stem rarely speckled, purple only at the nodes, scarcely glaucous, solid: leaves in 3's or 4's, rarely in 2's or 5's, sharply toothed: corolla 5-7 mm. long: heads paler than in the other species. Eastern Massachusetts and southern New Hampshire westward to Wisconsin, and southward to Pennsylvania, Kentucky, Oklahoma and Nebraska; also in the mountains from Virginia to Georgia. A plant of rich upland woods, rarely found near the coast.

The selection of names for these species has been found difficult. In the following discussion of the Linnean types the principle is observed that if there exists a specimen which Linnaeus had in hand when the description was written, this is to be considered the type, and is to be given the greatest weight in deciding upon the application of the name. If no such specimen exists, or if specimens and characters of two or more species are confused in the description, thus rendering definition by this means impossible, then the cited synonymy is to be employed in determining the type so far as it is of value.

In the first edition of the *Species Plantarum* (p. 838. 1753), Linnaeus gave the name *E. purpureum* to an American species of *Eupatorium* with verticillate leaves, and at the same time noted a var. β . In the *Amoenitates Academicæ*, *E. maculatum* was described by Linnaeus; and in the second edition of the *Species Plantarum* both species were treated at length. The citations under β in the

first edition all refer to our No. 1 above. Hermann's plate, and also that of Morrison, are plainly this plant; so also is the Hortus Cliffortianus plant in the British Museum as shown by a photograph. Moreover, the description accompanying the Ray citation represents this species, as shown by the number of leaves cited, as well as by a reference to the spotted stem and leaves like a nettle. Also, the original portion of the description of *E. purpureum* was drawn, either from the β references, or from the Hortus Cliffortianus plant, or both, and therefore refers to No. 1: "folia * * * lato-lanceolata * * * lanceolato-ovata * * * petiolata * * * Calyces florum incarnati. Flosculi octo."

In the Amoenitates, where *E. maculatum* was first proposed, all the citations under the original β are transferred to that species, and they are the only citations given. However, Linnaeus's description of *E. maculatum* applies much better to our No. 2: "foliis quinque tomentosis lanceolatis. * * * Folia quinque vel sex," since this northern plant usually has more lanceolate leaves, which are commonly more hairy underneath, and more frequently borne in 5's or 6's than is the case in plant No. 1. Plant No. 1 never has the leaves in 6's, and very rarely, if ever, in 5's. The situation is still further complicated by Linnaeus in the closing statement of the description where he says that *E. maculatum* is his variety of *E. purpureum* as to both synonymy and description. In the Linnean herbarium is a specimen from Kalm which Linnaeus must have had at the time the Species Plantarum was written, and which is undoubtedly plant No. 2. Moreover, plant No. 2, rather than No. 1, is much more likely to have been found by Kalm in the regions visited by him; since No. 1 is coastal, and not northern. A photograph of the specimen shows six leaves in the whorls (though unusual even for this species), and in every way answers the description of *E. maculatum* given by Linnaeus. There is every reason to suppose, therefore, that Linnaeus had this in hand when the description was drawn, and it is therefore to be considered the type-specimen. The name *E. maculatum* must, therefore, be applied to our No. 2, notwithstanding the fact that the synonymy applies to No. 1.

It remains to determine the application of the name *E. purpureum*. The statement has already been made that the original description seems to have been drawn from the β portion of the species. In the second edition of the Species Plantarum, where he first defines

together both *E. purpureum* and *E. maculatum*, Linnaeus recasts the description of *E. purpureum*, omitting the statement that the stem was spotted, and stating that it was, "viridis, ad exortum petiolarum purpurascens." This new description excludes both spotted-stemmed species, and narrows the application of the name to species Nos. 3 and 4. However, a still further study of the description shows that Linnaeus was confusing both of these, since the statement quoted above applies best to No. 4; while the statement "folio quina lanceolata-ovata. Calyces florum incarnati. Flosculi octo" applies to No. 3. A photograph of material now in the Linnaean herbarium seems to show, though an accurate identification of it is impossible, that Linnaeus at some time had specimens of both species. It is necessary, therefore, because of this confusion, to ignore the Linnaean description, and to attempt the definition of the name through the synonymy. The writer has not had access to the Colden citation. The plant on which the Gronovian reference was based was most probably our No. 3, as judged by the statement of Gronovius: "foliis ovato-lanceolatis obtuse serratis * * * foliis longis," and Gronovius's reference to Cornut which will later be shown to be No. 3. Moreover, though not much material has been seen from Virginia, it seems likely that No. 4 is largely confined to the mountains in that state, while No. 3 extends to the coast, and would be more likely to have been found by Clayton.

The next citation of Linnaeus was to Cornut where a plate and extended discussion is given. The plate is not convincing, as it might represent either species. In the text, however, is the statement: "Caules * * * rubescentes (cinereo tamen colore suffusi) [i. e., glaucous] * * * inanes intus," which can refer only to No. 3. The source of Cornut's plant, however, is in doubt, as our No. 3 is not known from Canada where the title of his work would lead one to infer that it was obtained.

The other citation of Linnaeus was to Morrison, but the figure there given seems to have been copied from Cornut, and the description is a verbatim transcription of Cornut's. The citations of Linnaeus, therefore, so far as they can be identified, unanimously refer to plant No. 3. Moreover, the key heading, "*Calycibus octo-floris," under which *E. purpureum* is found in the Species Plantarum, is appropriate to this plant and not to No. 4, which has from 3 to 6 florets. The name *E. purpureum* must therefore be used for our species No. 3.

In the *Species Plantarum*, on the page preceding that on which *E. purpureum* was described, Linnaeus proposed another species under the name *E. trifoliatum*. The original description consists solely of the statement "foliis ternis," which plainly is insufficient to define the species; and it is therefore necessary to depend upon the citations given, which are to Gronovius and Ray. The description given by Gronovius is inconclusive, having been drawn apparently from a slender shade plant with whitish heads without stating such characters as are of value here. The Clayton plant cited by Gronovius is his No. 620. A photograph of a specimen now in the British Museum labelled Clayton No. 620 shows it to be probably our No. 3. The leaves are lanceolate, bluntly and finely toothed; and, so far as can be made out from the print, the stem is purple and glaucous and not darker at the nodes. The stem is also cracked in one place in a manner more likely to occur if it were hollow. Also, as has already been stated, No. 3 is more likely to have been found by Clayton than No. 4. However, no species normally has leaves of this form in 3's. The specimen seems abnormal, but is more reasonably placed in No. 3. *E. trifoliatum* L. and *E. purpureum* L. are therefore to be considered one and the same species. Most of the early writers retained *E. trifoliatum* in addition to the various names employed by them for the other species. So far as the writer is aware, Torrey and Gray (Fl. N. A. ii. p. 82, 1841) were the first to unite *E. trifoliatum* and *E. purpureum*. This was done under the latter name; which, therefore, is the one to be retained under the Vienna Code (Chapter III, Sect. 5, Art. 46) which says: "When two or more groups of the same nature are united, the name of the oldest is retained. If the names are of the same date, the author chooses, and his choice cannot be modified by subsequent authors."

The oldest name that may be legitimately applied to our species No. 1 is the *E. verticillatum* of Lamarck (Encyc. ii, 405, 1786). This was divided by the author into two parts, α and β . Lamarck's citations and descriptions do not entirely agree. The Cornut and Morrison citations under α are the ones listed by Linnaeus under *E. purpureum* α , and have been shown above to refer to our No. 3. The Hermann reference under β , as has already been pointed out, refers to No. 1. However, the description given by Lamarck shows that both α and β are to be referred to species No. 1. In this description reference is made to a dull-green dotted stem and purple

heads with six or seven florets, a combination of characters appropriate to no other species than No. 1; and under β the statement that the stem is obscurely purple and spotted, leaves parallel-ribbed and not cottony, flowers about 8, must also refer to No. 1. It is true the statement under β of leaves in 5's is not appropriate to No. 1, which very rarely has 5 leaves in the whorl, and yet it could not refer to No. 2, for the number of florets is stated as only 8. Lamarck's description was drawn from a garden plant, and No. 1 might possibly have produced 5 leaves under cultivation. If greatest weight in determining the application of Lamarck's name is given to his description, then the name *E. verticillatum* must be applied to our species No. 1.

The oldest available name for the fourth species is apparently Michaux's *E. falcatum*. In his description the statement leaves 4-verticillate, oval-lanceolate, acuminate at each end and subfalcate, calyx 5-flowered, is most appropriate to No. 4. The panicle is stated as multicorymbose, and the flowers white. This species often has a very large panicle, and pale heads which are frequently nearly white in dense shade.

The following is a key to the species and varieties discussed:

- a. Leaves ovate to ovate-lanceolate, abruptly contracted into the petiole, more or less 3-nerved: plant somewhat viscid, scabrous-puberulent, with a strong odor when fresh: stem finely purple-speckled, not glaucous: inflorescence convex (leaves in 3's or 4's, very rarely in 2's or 5's: florets 6-9, rarely 5-12).....1. *E. verticillatum*.
 - a. Leaves lanceolate to oval, tapering at the base, pinnately veined (except sometimes in the variety of No. 2): plant not viscid and not odorous. b
 - b. Florets 9-15 (rarely 8-20): inflorescence or its divisions flat-topped: stem speckled if not obscured by too deep purple, not glaucous (leaves puberulent to glabrate beneath, in 4's or 5's, rarely in 3's or 6's: florets scarcely exserted). c.
 - c. Leaves below the inflorescence, and bracts, inconspicuous. 2. *E. maculatum*.
 - c. Leaves surrounding the inflorescence, and bracts, large and conspicuous, much exceeding the inflorescence, often more or less 3-nerved. var. *foliosum*.
 - b. Florets 5-7 (rarely 3-8): inflorescence convex: stems rarely speckled, more or less glaucous. d.
 - d. Stems fistulose, purple, plainly glaucous: leaves in 4's to 6's, bluntly toothed, scabrous-puberulent beneath or glabrate: florets scarcely exserted: corolla 3.5-4.8 mm. long, very rarely longer. 3. *E. purpureum*.
 - d. Stems solid, green with purple nodes, faintly glaucous: leaves in 3's or 4's, very rarely in 2's or 5's, sharply toothed, villous-pubescent beneath or glabrate: florets much exserted: corollas 5.5-7.5 mm. long: heads paler than in the other species.....4. *E. falcatum*.
1. *E. VERTICILLATUM* Lamarck, Encyc. ii. 405 (1786) α and β .
E. purpureum L. Sp. Pl. Ed. 1, ii. 838 (1753) as to synonymy under β , and as to description, but not as to synonymy under α , and not as

to his later treatment. *E. maculatum* L. Amoen. Acad. iv. 288 (1759) as to synonymy only. *E. fusco-rubrum* Walter, Fl. Car. 199 (1788) probably. *E. punctatum* Willd. Enum. Pl. Berol. 853 (1809) (photo. seen). *E. dubium* Willd. in Lam. Encyc. Sup. 2. 606 (1811). *E. ternifolium* Elliott, Bot. So. Car. and Ga. ii. 306 (1824) probably. *E. purpureum* var. *maculatum* Darlington, Fl. Cestrica 453 (1837).—Stem finely speckled with purple, green or more often suffused with purple, not glaucous, scarcely darker at the nodes, usually solid but frequently hollow, scabrous-puberulent above and apparently somewhat glandular: leaves generally in 3's or 4's occasionally in 2's or 5's, from broadly to narrowly ovate, sub-acuminate, abruptly narrowed to a conspicuous petiole which is 10–15 mm. long, coarsely serrate with somewhat rounded and mucronate or sharper teeth, veiny and often rugose, more or less plainly 3-nerved, above usually scabrous, beneath atomiferous and glabrous except on the veins which are more or less scabrous-puberulent: inflorescence rather small and dense, convex, when well developed hemispherical or short-oblong: heads narrowly oblong, 6–9 (rarely 5 or 12)-flowered, usually deep-purple: involucre bracts slightly narrower than in the next species; the inner often sub-acute: corolla 4.5–5.5 mm. long, slightly exerted: achenes 3.5–4.5 mm. long.—Borders of swamps and in marshes, in sandy or gravelly acid soils: on the Coastal Plain from eastern Massachusetts and southern New Hampshire to South Carolina. NEW HAMPSHIRE: Rochester, 1888, Mrs. E. Bartow; Derry, 1916, C. F. Batchelder; Mason, 1916, C. F. Batchelder; Jaffrey, 1896, W. Deane, 1897, B. L. Robinson, no. 379; Rindge, 1916, C. F. Batchelder; Gilsum, 1899, M. L. Fernald, no. 209. MASSACHUSETTS: eastern Massachusetts, 1854, W. Boott; North Andover, 1885, C. H. Morris; Wakefield, 1886, F. S. Collins; Sherborn, 1911, Martha L. Loomis, no. 430; South Royalston, 1907, J. A. Bates; Maugus Hill, Wellesley, 1897, E. F. Williams; Brookline, 1900, G. E. Morris (white flowers); Blue Hills, 1894, W. H. Manning; Cohasset, 1901, E. F. Williams; Loon Pond, Lakeville, 1913, Fernald & Long, no. 10490; Mashpee, 1916, Knowlton, Bean & Bird; East Sandwich, 1919, Fernald & Long, no. 19163; Allen's Harbor Creek Harwich, 1919, Fernald & Long, no. 19166; Brewster, 1912, F. S. Collins, no. 1649; Chilmack, 1911, J. A. Cushman, no. 7571; New Bedford, 1872, G. Mackie. RHODE ISLAND: Rumford, 1903, E. F. Williams; Providence, 1844, G. Thurber; Crescent Beach, Block Island, 1913, Fernald, Long & Torrey, no. 10491. CONNECTICUT: Waterford, 1899, C. B. Graves; Saybrook Junction, 1914, R. W. Woodward (flowers white); Southington, 1898, L. Andrews, no. 42, 1898, C. H. Bissell, no. 311; Tranquillity Farm, Middlebury, 1896, W. M. Shepardson. NEW YORK: Port Chester, 1886, Louise M. Stabler. NEW JERSEY: May's Landing, 1916, I. Tidestrom, no. 8042; Hammonton, 1917, A. Gershoy, no. 660; Dividing Creek, 1910, B. Long, no. 4843; Atsion, 1917,

A. Gershoy, no. 661; Bennett, 1917, *A. Gershoy*, no. 659. DELAWARE: near Delaware City, 1916, *I. Tidestrom*, no. 7903. DISTRICT OF COLUMBIA: Washington and vicinity, 1898, *E. S. Steele*. SOUTH CAROLINA: Santee Canal, *Ravenel*.

This is the commonest species in low ground along the Coastal Plain. Albino forms are occasionally found. The glandular pubescence apparently gives the plant a characteristic strong odor not present in the other species, but this observation needs verification. The stem is occasionally hollow. The variation in number of florets among the specimens studied was as follows: 4 with 5 florets, 17 with 6 florets, 13 with 7 florets, 10 with 8 florets, 11 with 9 florets, 5 with 10 florets, 2 with 11 florets, and 1 with 12 florets. The number of leaves in the whorl fluctuated in the following proportion: 4 with 2 leaves (small plants), 27 with 3 leaves, 28 with 4 leaves, and 1 with 5 leaves.

2. *E. MACULATUM* L. Amoen. Acad. iv. 288 (1759) as to original description and specimen in the Linnaean Herbarium, but not as to citations. *E. Bruneri* A. Gray, Synopt. Fl. i. pt. 2, 96 (1884). *E. atromontanum* A. Nelson, Bot. Gaz. xxxi. 400 (1901). *E. Rydbergii* Britton, Manual 921 (1901). *E. purpureum* var. *Bruneri* Robinson, Proc. Amer. Acad. xlii. 44 (1906) for the more hairy western plants.—Stem green and finely purple-speckled, more often the spots obscured by a deep purple suffusion, not glaucous, puberulent above, not darker at the nodes, very exceptionally hollow: leaves most commonly in 4's or 5's, rarely in 3's or 6's, elliptic-ovate or elliptic-lanceolate, tapering at base and apex, short petioled or nearly sessile, sharply and often irregularly incurved-serrate varying to more finely crenate-serrate, pinnately veined, rugose, above glabrous or slightly scabrous, beneath atomiferous and from nearly glabrous to canescent with minute crisp scabrous hairs on the veins or more commonly on both veins and parenchyma: inflorescence or its parts flat-topped, dense: heads broadly oblong, 9–15 (rarely 8 or 20)-flowered, usually deep purple: bracts of the involucre broad, obtuse: corollas 5 mm. long, their tips usually but slightly or not at all exerted beyond the involucre: achenes 3.4–4.2 mm. long.—Open grounds and the borders of thickets, along streams and in wet sedgy meadows, in rich often mucky scarcely sandy soils, generally in calcareous regions: Newfoundland and Quebec to northern New England, western Massachusetts, western Connecticut, and Lancaster County, Pennsylvania; westward through Michigan, Illinois, Colorado, Utah, and Saskatchewan to New Mexico and British Columbia. NEWFOUNDLAND: Salmonier River, 1894, *Robinson & Schrenk*, no. 40. QUEBEC: Grindstone Island, Magdalen Islands, 1912, *Fernald, Long & St. John*, no. 8093; vicinity of Cap a L'Aigle, 1905, *John Macoun*, no.

68338. PRINCE EDWARD ISLAND: Hillsborough River, Mt. Stewart, 1912, *Fernald, Bartram, Long & St. John*, no. 8092; Brackley Point, 1912, *Fernald, Long & St. John*, no. 8094. MAINE: Lake Hadley, East Machias, 1898, *M. A. Barber*; Orono, 1887, *M. L. Fernald*, 1896, *E. D. Merrill*, no. 437; Dover, 1895, *M. L. Fernald*, no. 307; Abbott, 1916, *Fernald & Long*, no. 14647; Fairfield, 1916, *Fernald & Long*, no. 14646; Greenvale, 1894, *K. Furbish*; South Deer Isle, 1914, *A. F. Hill*, no. 1834; Cow Island, Topsham, 1910, *K. Furbish*; South Poland, 1895, *K. Furbish*; Wells, 1898, *K. Furbish*. NEW HAMPSHIRE: Shelburne, 1915, *W. Deane*; Dartmouth College Grant, 1914, *A. S. Pease*, no. 16194; Gate of Crawford's Notch, 1884, *C. E. Faxon* (albino form); Walpole, 1916, *C. F. Batchelder*. VERMONT: Willoughby, 1898, *G. G. Kennedy*; Ripton, 1908, *E. F. Williams*; Manchester, 1898, *M. A. Day*, no. 91. MASSACHUSETTS: Greenfield, 1913, *J. Murdoch, Jr.*, no. 5208; Worthington, 1912, *B. L. Robinson*, no. 596; Cheshire and vicinity, 1912, *E. J. Winslow*; Lenox, 1911, *R. Hoffmann* (shade form); Stockbridge, 1902, *R. Hoffmann*; New Marlboro, 1912, *R. Hoffmann*; Mount Washington, 1915, *F. G. Floyd*. NEW YORK: Canton, 1914, *Orra P. Phelps*; Ithaca, 1913, *E. L. Palmer*, nos. 1190 & 1191; 1915, *C. C. Thomas*, no. 5099; 1919, *A. J. Eames*, nos. 12973, 12975, 12978, 12979; 1919, *Eames & Wiegand*, no. 12974; 1919, *Wiegand*, no. 12976; 1919, *L. F. Randolph*, no. 12972; Big Gully Ravine, Springport, 1919, *Eames, Randolph & Wiegand*, no. 12980; North Spencer, 1915, *Eames & MacDaniels*, no. 5088; Spencer Lake, 1919, *Eames & Wiegand*, no. 12981. PENNSYLVANIA: Dillerville Swamp, in limestone, 1901, *A. A. Heller*. ONTARIO: Plevna, 1902, *J. Fowler*; Island Lake, Algonquin Park, 1900, *Macoun*, no. 21813. OHIO: Birmingham, 1914, *L. H. MacDaniels*, no. 151. MICHIGAN: Keweenaw County, 1890, *O. A. Farwell*, no. 777; Turin, 1901, *B. Barlow*; Flint, 1909, *E. E. Sherff*; South Haven, 1911, *O. E. Lansing, Jr.*, no. 3316. INDIANA: East Chicago, 1910, *O. E. Lansing, Jr.*, no. 2804. ILLINOIS: Ravinia, 1911, *E. E. Sherff*; Cedar Lake, Lake County, 1906, *Gleason & Shobe*, no. 142; South Chicago, 1913, *H. H. Smith*, no. 5752; near Wady Petra, 1900, *V. H. Chase*, no. 764. WISCONSIN: Fort Howard Marsh, Brown County, 1890, *J. H. Schuette*; Osceola, 1900, *C. F. Baker*. IOWA: Ames, 1896, *Pammel & Ball*, no. 34. MINNESOTA: Cass Lake, Cass County, 1914, *L. H. & H. E. Pammel*, no. 553; Muskoda, Red River Valley, 1901, *C. A. Ballard*, no. 3083. NORTH DAKOTA: near Bottineau, 1902, *J. Lunell*. SOUTH DAKOTA: Black Hills, 1910, *J. Murdoch, Jr.*, no. 4296. NEBRASKA: sand hills of central Nebraska, 1893, *P. A. Rydberg*, no. 1682; Fort Niobrara, Knox County, 1908, *J. M. Bates*, no. 4686; Paddock, 1893, *F. Clements*, no. 2797. COLORADO: Fort Collins, 1881, *Bruner* (type of *E. Bruneri* Gray). WYOMING: Beaver Creek, Black Hills, 1896, *A. Nelson*, no. 2553 (type number of *E. atromontanum* Nelson). UTAH: Wasatch Mountains, 1872,

Wheeler. NEW MEXICO: Fort Wingate, *Dr. Matthews*. SASKATCHEWAN: 1857-58, *E. Bourgeau*. BRITISH COLUMBIA: Fraser River, 1858, *Lyall*; Chilliwack Valley, 1901, *J. M. Macoun*, no. 26486.

Var. **foliosum** (Fernald) comb. nov. *E. purpureum* var. *foliosum* Fernald, RHODORA x. 86 (1908).—Leaves large, ovate-oval, with a tendency toward a 3-ribbed base, thin, nearly or quite glabrous: inflorescence small, surrounded and usually much surpassed by the large lance-ovate upper leaves and bracts.—Newfoundland and Saguenay County, Quebec to northern New England and possibly Lake Superior. NEWFOUNDLAND: Manuel's River, 1894, *Robinson & Schrenk*; Brigus Junction, 1911, *Fernald & Wiegand*, no. 6275; Grand Falls, 1911, *Fernald & Wiegand*, no. 6276; Bluff Head, Bay of Islands, 1898, *A. C. Waghorne*, no. 18; Harry's River, 1910, *Fernald & Wiegand*, no. 4133. QUEBEC: Dartmouth River, 1904, *Collins, Fernald & Pease*; Mingan, 1915, *H. St. John*, no. 90749. PRINCE EDWARD ISLAND: Bloomfield, 1912, *Fernald, Long & St. John*, no. 8095. NEW BRUNSWICK: Kent County, 1870, *J. Fowler*. MAINE: Van Buren, "the common form," 1900, *M. L. Fernald*; Hamlin, 1901, *Fernald*; Fort Fairfield, 1901, *E. F. Williams*; Boundary Lake, 1902, *Eggleston & Fernald*; Fairfield, 1916, *Fernald & Long*, no. 14648; Bridgton, 1912, *L. R. Martin*. NEW HAMPSHIRE: Ice Gulch, Randolph, 1908, *A. S. Pease*, no. 11498; Pittsburgh, *Fernald & Pease*, no. 17068; Jefferson, 1916, *Pease*, no. 16837. MICHIGAN: moist thickets, Keweenaw County, 1887, *O. A. Farwell*, no. 441 (doubtful).

E. maculatum is a northern plant of the richer, more calcareous soils, and is therefore nearly or quite absent from the sandy coastal plain from New England southward. In Central New York it is the characteristic joe-pye weed, occurring in great abundance everywhere in low grounds except perhaps in acid bogs. Locally it is somewhat variable as affected by shade and moisture, but geographically it is much more so. In the northeast the var. *foliosum* seems distinct enough to warrant separate treatment, but the other variations pass one into the other so gradually that no practical dividing lines can be drawn. Thus the leaves of the northeastern specimens are only slightly hairy beneath, but as one travels westward they gradually become more hairy until some plants on the Great Plains have the leaves distinctly whitened on the under surface. Locally, too, the degree of pubescence may fluctuate with the exposure. The more canescent western plants have been separated by Robinson as *E. purpureum* var. *Bruneri* (Gray) Robins. (l. c.). Several sheets of var. *foliosum* from Gaspé have the upper portion of the stem, and to a certain extent the under surface of the leaves, villous instead of

crisp-puberulent. Scattered specimens of the typical form have narrow leaves with fine crenate serrations resembling those of *E. purpureum*, a variation the significance of which is not understood but which is probably ecological. The florets vary considerably in number, due apparently in some cases at least to local conditions. Among the specimens examined the number of florets ran as follows: 1 with 7 florets, 2 with 8 florets, 10 with 9 florets, 12 with 10 florets, 12 with 11 florets, 30 with 12 florets, 17 with 13 florets, 10 with 14 florets, 7 with 15 florets, 1 with 16 florets, 3 with 17 florets, 1 with 18 florets, 1 with 19 florets, 0 with 20 florets, and 1 with 21 florets. The leaves in the whorls fluctuate within narrow limits. The number counted ran as follows: 1 with 2 leaves (starved specimen), 7 with 3 leaves, 55 with 4 leaves, 54 with 5 leaves, and 2 with 6 leaves.

3. *E. PURPUREUM* L. Sp. Pl. ed. 1, ii. 838 (1753) as to synonymy under α , which in this case determines the type. *E. trifoliatum* L. Sp. Pl. ed. 1, ii. 837 (1753), placed under *E. purpureum* by T. & G. in 1841. *E. americanum* Hill, Brit. Herb. 453 (1756). ? *E. purpureum* γ *angustifolium* T. & G. Fl. N. Amer. ii. 82 (1841). *E. fistulosum* Barrett, Eupat. verticillate (1841), see also Wood's Class Book, ed. 2, 314 (1847).—Stems normally unspeckled, occasionally mottled, uniformly suffused with purple and not darker at the nodes, plainly glaucous, glabrous, hollow: leaves in 4's–6's, very rarely in 7's, elliptic-lanceolate, acuminate, tapering gradually to the nearly sessile base, regularly and finely crenate-serrate, scarcely rugose, above glabrous, beneath atomiferous and sparingly fine-puberulent on the veins only, or commonly almost glabrous; the veins numerous, spreading, regularly decreasing toward base and apex of the leaf; inflorescence crisp-pubescent, convex, often hemispherical, when well developed large and loose with the lower branches divaricate or horizontal: heads purple, narrowly cylindrical, 6–7 (rarely 5 or 8)-flowered: bracts of the involucre mostly obtuse: corolla 3.5–4.8 mm. long, slightly or not at all exserted: achenes 3.2–4.5 mm. long.—Low fields and the borders of thickets, if not too wet, in rich sandy and gravelly scarcely calcareous soils: southern Maine to Rhode Island, Florida, Texas and Oklahoma, and from eastern Kentucky, and West Virginia through western Pennsylvania to Ohio. MAINE: Cape Elizabeth, 1911, *M. L. Fernald*; Biddeford Pool, 1900, *G. G. Kennedy*. MASSACHUSETTS: Concord, *E. S. Hoar*; Milton, 1899, *G. G. Kennedy*; East Sandwich, 1919, *Fernald & Long*, no. 19168; Deerfield, 1908, *M. A. Day*, no. 62. RHODE ISLAND: Providence, *G. Thurber*. CONNECTICUT: Pomfret, 1910, *Sarah R. Armington*; Gully Brook, Hartford, 1907, *A. W. Driggs*; Ledyard, 1901, *C. B. Graves*, no. 234. MARYLAND: Ellicott City, 1916, *G.*

Arsène. DISTRICT OF COLUMBIA: 1901, *E. S. Steele*; Brookland, 1908, *T. Holm*; near Fort Totten, 1915, *T. Holm*. VIRGINIA: Rapahannock River, 1915, *I. Tidestrom*, no. 7607; Bedford County, 1871 & 1872, *A. H. Curtiss*. WEST VIRGINIA: Harman, 1904, *A. H. Moore*, no. 2167; Dry Fork River near Harman, 1904, *J. M. Greenman*, no. 236. KENTUCKY: near Poor Fork Post Office, 1893, *T. H. Kearney, Jr.*, no. 214. SOUTH CAROLINA: Santee Canal, *H. W. Ravenel*. ALABAMA: *Buckley*. FLORIDA: Eustis, 1895, *G. V. Nash*, no. 2118. TEXAS: Pope, *ex herb. G. Thurber*. OKLAHOMA: Page, 1914, *O. W. Blakley*, no. 3417. ARKANSAS: southwestern Arkansas, *F. L. Harvey*, no. 4. PENNSYLVANIA: Pittsburgh, 1831, *Holz*. OHIO: Little Mountain, 1897, *J. M. Greenman*, no. 310.

In general this plant is very constant in its characters. One specimen from Pittsburgh, Pa. (*Holz*) has corollas longer than 5 mm.; otherwise this character of the corolla is good. The variation in florets was: 6 with 5 florets, 14 with 6 florets, 5 with 7 florets, and 1 with 8 florets. The variation in number of leaves in the whorl was: 7 with 4 leaves, 5 with 5 leaves, 6 with 7 leaves, and 1 with 8 leaves.

- F. trilobatum L. *see Rhod. 29; 6. 1927.*

4. *E. FALCATUM* Michx. Flor. Bor. Am. ii. 99 (1803). *E. purpureum* L. Sp. Pl. ed. 2, 1173 (1763) as to description in part, not as to synonymy. *E. amoenum* Pursh, Fl. Am. Sept. ii, 514 (1814). *E. purpureum* var. *amoenum* A. Gray, Synopt. Fl. N. A. i. pt. 2, 96 (1884). *E. purpureum* var. *falcatum* Britton, Mem. Torr. Bot. Club v. 312 (1894).—Stems normally unspeckled, rarely mottled, green, purple at the nodes, rarely more purplish, scarcely glaucous, glabrous or nearly so, solid: leaves mostly in 4's, less commonly in 3's, rarely in 2's or 5's, lanceolate to ovate-oval or rarely ovate, acuminate, tapering gradually to the petioled or nearly sessile base or rarely more abruptly tapering, sharply serrate, slightly rugose; veins numerous, less spreading and less regular than in the last, decreasing gradually toward base and apex; blade glabrous and nearly smooth above, atomiferous and from glabrous to densely crisp-villous beneath, the hairs longer and softer than in the other species: inflorescence crisp-downy, convex, when well developed hemispherical or short-oblong and loose, often very large and open (up to 5 dm. long and 4 dm. wide): heads narrowly cylindrical, pale purple or whitish, 3–6 (7)-flowered: involucrel bracts narrowly oblong, the inner acutish: corollas (5-) 5.5–7.5 mm. long, much exerted beyond the involucre: achenes 3–5 mm. long.—Open woods and wood borders in damp or rather dry rich light soil, but not in the coastal sands: eastern Massachusetts, southern New Hampshire, and southern Vermont, to Connecticut, and in the uplands to Georgia; westward through Ontario, New York, and Pennsylvania to Wisconsin, Nebraska and Oklahoma.

NEW HAMPSHIRE: Asquam Lake, 1914, *A. L. Gundersen*; Walpole, 1899, *M. L. Fernald*, no. 214. VERMONT: Pownal, 1898, *J. R. Churchill*, also *W. W. Eggleston*, no. 258; 1899, *Eggleston*, no. 1. MASSACHUSETTS: Revere, 1879 & 1882, *H. A. Young*; Middlesex Fells, 1894, *W. H. Manning*; Blue Hills, 1894, *C. G. French*; Readville, *C. E. Faxon*; Stockbridge, 1902, *R. Hoffmann*. CONNECTICUT: Waterbury, 1910, *A. E. Blewitt*, no. 1135; Hammond's Woods, Waterford, 1902, *C. B. Graves*, no. 302; New Haven, *D. C. Eaton*. NEW YORK: Oneida Lake, *old Torrey & Gray specimen*; various places in the Cayuga Lake Basin, 1919, *Eames, Randolph & Wiegand*, nos. 12983, 12984, 12985, 12986, also *Eames & Wiegand*, nos. 12982, 12989, and *A. J. Eames*, no. 12987; Junius, 1915, *Eames & MacDaniels*, no. 5090. VIRGINIA: Stony Man Mountain near Luray, 1901, *E. S. & Mrs. Steele*, no. 9; Bedford County, 1871 & 1872, *A. H. Curtiss*. WEST VIRGINIA: Parsons, 1904, *A. H. Moore*, no. 1994; East Fork of the Greenbrier River, Pocohontas County, 1904, *A. H. Moore*, no. 2380, and *J. M. Greenman*, no. 235; Gap Mountain, 1903, *E. S. & Mrs. Steele*, no. 186. KENTUCKY: on Big Black Mountain, Harlan County, 1893, *T. H. Kearney, Jr.*, no. 166. NORTH CAROLINA: Swain County, 1891, *Beardslee & Kofoid*. GEORGIA: north Georgia, 1875, *C. Wright*; Whitfield County, 1900, *P. Wilson*, no. 70; west of Cuthbert, 1903, *R. M. Harper*, no. 1877. ONTARIO: Queenston, 1911, *J. White*, no. 2. MICHIGAN: near Port Huron, 1892, *C. K. Dodge*; Flint, 1909, *E. E. Sherff*. INDIANA: Wabash River east of Bluffton, 1908, *C. C. Deam*, no. 5194. WISCONSIN: Brown County, 1901, *J. H. Schuette* (albino form); Kankauna, 1890, *Schuette*. ILLINOIS: Bloomington, 1886, *B. L. Robinson*; near Princeville, 1900, *V. H. Chase*, no. 716; Peoria, 1904, *F. E. MacDonald*; Havana, 1903, *H. A. Gleason*; St. Clair County, 1886, *H. Eggert*; Grand Tower, 1902, *H. A. Gleason*; Belknap, 1902, *Gleason*. IOWA: Ames, 1897, *Ball & Pammel*; Dakota City, 1896, *Pammel*, no. 33. NEBRASKA: Nemaha, 1910, *J. M. Bates*, no. 5211. OKLAHOMA: Grand River, Cherokee Nation, 1895, *J. W. Blankinship*; near Ottawa, 1913, *G. W. Stevens*, no. 2371.

E. falcatum is much more variable than *E. purpureum*. In the southern mountains it is often very slender (*E. amoenum* Pursh) but ordinarily it is large and frequently very tall with an expanded inflorescence as noted in the description. The nodes occasionally lose the purple coloration. The leaves vary widely in shape, and the pubescence is extremely variable in density, but it is always of the same loose type. The heads are usually pale, but may vary to white or to deeper purple. The transitions in all cases are so gradual, and the combinations so complex, that no subdivision of the species has seemed practical. The variation in florets was as follows: 4

with 3 florets, 7 with 4 florets, 25 with 5 florets, 24 with 6 florets, and 2 with 7 florets. The variation in number of leaves in a whorl was: 2 with 2 leaves (poor specimens), 20 with 3 leaves, 36 with 4 leaves, and 2 with 5 leaves.

CORNELL UNIVERSITY, Ithaca, New York.

THE AMERICAN AMMOPHILA.

M. L. FERNALD.

THE common Sand Reed, Psamma, Marram or Beach Grass, which covers the coastal sand dunes from the Straits of Belle Isle to North Carolina and occurs on sandy shores of the St. Lawrence system inland quite to Lake Superior, has been universally identified with *Ammophila arenaria* (L.) Link, the species occurring on the western and southern coasts of Europe. Superficially the two are very similar, although it needs only a glance at good material of typical *A. arenaria*, which occurs from southern Scandinavia to Portugal and Morocco, to see that the spike-like panicle is much shorter than in most of the American plant, *A. arenaria* having panicles only 0.5-2 dm. long, the panicles of the Atlantic American plant ranging from 1.5-4 dm. in length. In its long panicle the American plant is more nearly approached by the Mediterranean *A. arenaria*, var. *arundinacea* (Host) Husnot, in which the panicles may be 3 dm. long.

In all its technical characters, however, the long-panicled Mediterranean *Ammophila arenaria*, var. *arundinacea* agrees with the more northwestern typical European *A. arenaria*; but in these characters the European and Atlantic American plants are quite distinct, the Old World species being known in America only on the Pacific coast, where it has very recently been introduced as a sand binder. Briefly stated, the two species differ as follows: In *A. arenaria* the upper surface of the leaf-blade is copiously puberulent along the cartilaginous nerves, in the American merely serrulate-scabrous; in *A. arenaria* the ligule is scarious, lance-attenuate and very prolonged, commonly 1.5-3 cm. long, and lacerate at tip; in the Atlantic American plant, on the other hand, the ligule is chartaceous or coriaceous, rounded and very short, ranging from 1-3 mm. in length. In *A. arenaria*,

the lemma is sharply bidentate at tip and short-mucronate in the sinus and the palea is sharp-mucronate; but in the American plant the lemma and palea are both blunt, the former scarcely bidentate. Other differences of less taxonomic importance are the puberulent glumes of the American species contrasted with the more commonly glabrous glumes of the European; and, similarly, the puberulent axis of the panicle in the American, the axis in the European being nearly or quite glabrous.

That the two are thoroughly distinct species there can be no question; but the American plant seems to have no valid name. In 1818, Nuttall published the name *Phalaris maritima*¹ partly for a plant of New Jersey, but he gave absolutely no description and stated that his plant was *Arundo arenaria* L. and that it is an important grass of Europe. It would be entirely unwise to take up for our grass Nuttall's name which was intended merely as a substitute for *Arundo arenaria*. The Atlantic American plant may be called

AMMOPHILA breviligulata, n. sp., *A. arenariam* simulans; foliis supra serrulato-scabris, ligulis chartaceis vel coriaceis rotundatis 1–3 mm. longis; paniculis lineari-cylindricis 1.5–4 dm. longis, rhachi puberulo; glumis puberulis; lemmate obtuso nec bidentato; palea obtusa; caryopsibus 3.2–3.6 mm. longis.—Sand dunes and shores, Newfoundland to North Carolina, inland along the St. Lawrence system to the Great Lakes. TYPE: sandy sea-beach, Milford, Connecticut, August 27, 1902, *C. H. Bissell* (Gray Herb.).

GRAY HERBARIUM.

¹ Nutt. Gen. i. 48 (1818).

REPORTS ON THE FLORA OF THE BOSTON
DISTRICT,—XXXII.

VITACEAE.

CISSUS.

C. AMPELOPSIS Pers. Ipswich, growing in a dense mass over boulder in wild ground, perhaps from a neighboring cemetery (*R. A. Ware*, Oct. 16, 1909).

PSEDERA.

P. quinquefolia (L.) Greene. Dry rocky soil, common.

P. quinquefolia (L.) Greene, var. *hirsuta* (Donn) Rehder. Dry rocky woods, Boxboro (*F. W. Hunnewell & H. St. John*, May 30, 1914).

P. vitacea (Knerr) Greene. Medford, Concord, Wellesley, Sherborn.

VITIS.

V. aestivalis Michx. Dry open woods and thickets, apparently frequent, except in the southeast.

V. bicolor Le Conte. Woods and thickets, common.

V. labrusca L. Woods and roadsides, common throughout.

V. novae-angliae Fernald. Georgetown (*E. F. Williams*, Aug. 9, 1907); Sudbury (*C. W. Swan*, Sept. 2, 1885). See RHODORA xix. 144, 1917.

V. vulpina L. Sandy soil at Gloucester, Essex, Beverley, Ipswich and Bradford.

TILIACEAE.

TILIA.

T. americana L. Rich woods, frequent.

MALVACEAE.

ABUTILON.

A. THEOPHRASTI Medic. Gardens and waste places, frequent, especially near Boston.

ALTHEA.

A. OFFICINALIS L. A single plant on rubbish heap in Cambridge (*W. Deane*, Aug. 5, Oct. 3, 16, 1886); reported from marsh at Salisbury according to J. Robinson, Fl. Essex Co. 40, 1880.

ANODA.

A. TRIANGULARIS (Willd.) DC. Woolwaste dump, Westford (*Miss E. F. Fletcher*). See RHODORA xviii. 143, 1916. Native of southwestern United States.

GOSSYPIUM.

G. HERBACEUM L. Self-sown and abundant on vacant lot on Huron Ave., Cambridge (*E. F. Williams*, Sept. 12, 1910). Probably an Asian plant.

HIBISCUS.

H. Moscheutos L. Salt or brackish marshes; Salisbury (*Mrs. C. N. S. Horner*, —, 1882); old report at Swampscott (J. Robinson, Fl. Essex Co. 40, 1880); frequent from Hingham south. There is an especially fine station nearly an acre in extent at Mann's Hill, Scituate. This fine mallow also grows in fresh water marshes at various places along the Concord and Sudbury rivers as far inland as Framingham; there are occasional stations on the Charles River, and the plant is rather abundant at the northeast of Farm Pond, Sherborn (*A. J. Eames*).

H. TRIONUM L. Gardens and waste places, occasional.

MALVA.

M. ALCEA L. Escaped from gardens, or persistent, at seven places.

M. CRISPA L. Casual at Reading, Malden, S. Boston, Mansfield and Ashland. Native of Europe, and sparingly escaped from old gardens and elsewhere.

M. MOSCHATA L. Persistent and spreading.

M. PARVIFLORA L. Rubbish heap, Cambridge (*W. Deane*, Oct. 11, 1884). Introduced probably from the Pacific coast, where it is well established.

M. PUSILLA Smith (*M. borealis* Wallm. of Middlesex Flora). Vicinity of woollen mills at Westford (*C. W. Swan*, Sept. 15, 1884); Dracut (*C. W. Swan*, Aug. 1, 1883); also Lowell (*C. W. Swan*, no date). Native of northern Europe and Asia, introduced into California, and casual here. See Synop. Fl. N. Amer., A. Gray, ed. B. L. Robinson, i. pt. 1, 298-299, 1897.

M. ROTUNDIFOLIA L. Gardens and waste places, common.

M. SYLVESTRIS L. Sharon (*S. Harris*, July 12, 1894). Reported from dry open roadsides, Acton and Concord, Hosmer, RHODORA i. 223, 1899.

M. SYLVESTRIS L., var. *MAURITIANA* (L.) Boiss. Charlestown (*C. E. Perkins*, July 23 and Sept. 9, 1881). Specimens in herb. N. E. Botanical Club and Gray Herb. Casual from southwestern Europe.

MALVASTRUM.

M. COROMANDELIANUM (L.) Garcke. Woolwaste, Westford, Miss E. F. Fletcher in RHODORA xix. 132, 1917. Introduced from India.

SIDA.

S. HERMAPHRODITA (L.) Rusby. Escaped from Fenway, Boston, also in field at West Cambridge. Persisting about old gardens, Jamaica Plain (*N. T. Kidder*, Sept. 20, 1888).

S. SPINOSA L. Dumps at Lowell, Dracut, Westford, Malden and Watertown.

SPHAERALCEA.

S. MUNROANA (Dougl.) Spach. Woolwaste dump, Westford, 1915, Miss E. F. Fletcher in RHODORA xviii. 143, 1916. Native of Rocky Mts. and portions of the northwest.

S. FENDLERI Gray. Woolwaste, Westford, Miss E. F. Fletcher in RHODORA xx. 20, 1918. Native of western Texas and Arizona.

WISSADULA.

W. CALLIMORPHIA (Hochr.) Hassl., var. *FRIESII* Hassl. Woolwaste, Westford (*Miss E. F. Fletcher*, Sept. 5, 1917). Native of eastern Bolivia and adjacent Brazil. See RHODORA xx. 20, 1918.

TAMARICACEAE.**TAMARIX.**

T. GALLICA L. Vacant lot, escaped from Fenway, Boston (*E. F. Williams*, May 23, 1909).

T. PARVIFLORA DC. Waste lands in Boston and Brookline, escaped from cultivation.

HYPERICACEAE.**HYPERICUM.**

H. AUREUM Bartram. Roadside, Billerica and Westford (*Miss E. F. Fletcher*, August, 1909). See *RHODORA* xii. 55, 1910.

H. boreale (Britton) Bicknell. Shores of ponds and wet places, common.

H. canadense L. Moist or dry soil, very common throughout.

H. ellipticum Hook. Wet places, frequent northward, local southward.

H. gentianoides (L.) BSP. Dry sand and gravel, also on ledges; common throughout.

H. majus (Gray) Britton. Wet or dry soil, common.

H. mutilum L. Moist soil, common throughout.

H. PERFORATUM L. Fields and pastures, especially in dry soil, very common throughout.

H. PROLIFICUM L. Weston (*E. J. Winslow*, July 9, 1911, et seq.).

H. punctatum Lam. Open places in damp and dry soil, frequent.

H. virginicum L. Swamps and wet shores, common throughout.

ELATINACEAE.**ELATINE.**

E. minima (Nutt.) Fisch. & Mey. Wet sandy margins of ponds, frequent. See *RHODORA* xix. 10-15, 1917.

C. H. KNOWLTON	} Committee on
WALTER DEANE	

JUNCUS GERARDI Loisel., var. **pedicellatus**, n. var., cum inflorescentia 1-2 dm. longa; floribus plerumque pedicellatis, pedicellis 3-10 mm. longis; perianthiis 3.5-5 mm. longis.

Inflorescence 1-2 dm. long: flowers mostly pedicelled; the pedicels 3-10 mm. long: perianths 3.5-5 mm. long.—Wet sandy, gravelly or rocky coast, Maine to Rhode Island. MAINE: Cutler, July 24, 1902, *Kate Furbish*; Dark Harbor, Islesboro, August 14, 1913, *Woodward, Bissell & Fernald*, no. 9,142; Matinicus, September, 1908, *C. A. E. Long*, no. 38 (TYPE in herb. N. E. B. C.); August 9, 1919, *Long*, no. 147. RHODE ISLAND: East Providence, July 20, 1890, *J. F. Collins*.

Differing from typical *J. Gerardi* in its large flowers and elongated pedicels, the perianths of the typical form of the species being 2-3 (rarely 3.5) mm. long, and the flowers mostly sessile or merely short-pedicelled. The extreme material of var. *pedicellatus* is very large, nearly 1 m. high, but other plants are low (about 3.5 dm.). The inflorescence, too, is longer than in most *J. Gerardi*, in which the inflorescence varies from 0.1-1 (rarely to 1.5) dm. long.—M. L. FERNALD, Gray Herbarium.

Vol 22, no. 255, including pages 33 to 56, was issued 20 April, 1920.

DUPLICATE BOOKS FOR SALE.

Kerner von Marilaun, A. & Oliver, F. W.	
The natural history of plants. 4 vols. 1895.	
New York. 1. 8°.....	\$10.00
Presents fascinatingly the relations of plants to their environment.	
Masters, M. T. Vegetable teratology. London.	
1869. 8°.....	\$6.00
The one standard work in English on this subject.	
Pickering, Charles. Chronological history of	
plants. Boston, 1879. 4°.	\$15.00
Spruce, R. Notes of a botanist on the Amazon	
& Andes. 2 vols. London, 1908. 8°.....	\$3.00
Sullivant, W. S. Icones Muscorum, 8vo., illus-	
trated by numerous copper plates of high ex-	
cellence (unbound and in sheets). Here offered	
at a greatly reduced price.....	\$6.00
Supplement to the preceding (also in sheets).	\$5.00
Icones and Supplement together.....	\$10.00
United States Exploring Expedition during 1838-	
1842, under command of Charles Wilkes. Un-	
bound, in sheets. Botany. Phanerogamia	
Part I, by A. Gray, with atlas. 1854.....	\$40.00
Watson, S. Bibliographical Index to N. A. Botany.	
Part I (all published). Washington. 1878. 8°	\$1.25
Address Librarian, GRAY HERBARIUM of HARVARD UNIVERSITY, Cambridge, Mass.	

INDICATORS FOR DETERMINING REACTIONS OF SOILS

As used by DR. EDGAR T. WHERRY in the studies described on page 35.
A set of six indicators, made up ready for use, in box suitable for carrying into the field. Accompanied by instructions for using them. Price \$2.85. For sale by: LAMOTTE CHEMICAL PRODUCTS CO., 13 W. Saratoga St., Baltimore, Md.

Driers Presses

Collecting Picks

Mounting Papers

Collecting Cases

Genus Covers

Herbarium Supplies

CAT. 91,—BOTANICAL. CAT. 92,—BIOLOGICAL, SLIDES, ETC.

Microscopes

Accessories

Lenses

CAMBRIDGE BOTANICAL SUPPLY COMPANY

Laboratory

WAVERLEY, MASS,

Equipment

8 Lexington Street